



BE PREPARED FOR A SPORTS INJURY!

The term “sports injury,” in the broadest sense, refers to the kinds of injuries that most commonly occur during sports or exercise. Some sports injuries result from accidents; others are due to poor training practices, improper equipment, lack of conditioning, or insufficient warm-up and stretching.

Common sports injuries can include:

- Muscle sprains and strains
- Tears of the ligaments that hold joints together
- Tears of the tendons that support joints and allow them to move
- Dislocated joints
- Fractured bones, including vertebrae

Whether an injury is acute or chronic, there is never a good reason to try to “work through” the pain of an injury. When you have pain from a particular movement or activity, STOP! Continuing the activity only causes further harm.

WHEN TO SEEK MEDICAL TREATMENT

You should call a health professional if:

- The injury causes severe pain, swelling, or numbness.
- You can't tolerate any weight on the area.
- The pain or dull ache of an old injury is accompanied by increased swelling or joint abnormality or instability.

WHEN AND HOW TO TREAT AT HOME


If you don't have any of the above symptoms, it's probably safe to treat the injury at home—at least at first. If pain or other symptoms worsen, it's best to check with your health care provider. Use the **RICE** method to relieve pain and inflammation and speed healing. Follow these four steps immediately after injury and continue for at least 48 hours.

REST Reduce regular exercise or activities of daily living as needed. If you cannot put weight on an ankle or knee, crutches may help. If you use a cane or one crutch for an ankle injury, use it on the uninjured side to help you lean away and relieve weight on the injured ankle.

ICE Apply an ice pack to the injured area for 20 minutes at a time, four to eight times a day. A cold pack, ice bag, or plastic bag filled with crushed ice and wrapped in a towel can be used. To avoid cold injury and frostbite, do not apply the ice for more than 20 minutes. (Note: Do not use heat immediately after an injury. This tends to increase internal bleeding or swelling. Heat can be used later on to relieve muscle tension and promote relaxation.)

COMPRESSION Compression of the injured area may help reduce swelling. Compression can be achieved with elastic wraps, special boots, air casts, and splints. Ask your health care provider for advice on which one to use.

ELEVATION If possible, keep the injured ankle, knee, elbow, or wrist elevated on a pillow, above the level of the heart, to help decrease swelling.



Rest. Ice. Compression. Elevation.

THE BODY'S HEALING PROCESS

From the moment a bone breaks or a ligament tears, your body goes to work to repair the damage. **Here's what happens at each stage of the healing process:**

At the moment of injury: Chemicals are released from damaged cells, triggering a process called inflammation. Blood vessels at the injury site become dilated; blood flow increases to carry nutrients to the site of tissue damage.

Within hours of injury: White blood cells (leukocytes) travel down the bloodstream to the injury site where they

begin to tear down and remove damaged tissue, allowing other specialized cells to start developing scar tissue.

Within days of injury: Scar tissue is formed on the skin or inside the body. The amount of scarring may be proportional to the amount of swelling, inflammation, or bleeding within. In the next few weeks, the damaged area will regain a great deal of strength as scar tissue continues to form.

Within a month of injury: Scar tissue may start to shrink, bringing damaged, torn, or separated tissues back together. However, it may be several months or more before the injury is completely healed.

WHO SHOULD I SEE FOR MY INJURY?

Although severe injuries will need to be seen immediately in an emergency room, particularly if they occur on the weekend or after office hours, most musculoskeletal sports injuries can be evaluated and, in many cases, treated by your primary health care provider.

